

Gumbo Limbo Spiraling Whitefly A New Whitefly in South Florida



Adult whitefly

Introduction: In March, 2009, a whitefly (*Aleurodicus rugioperculatus* Martin: Hemiptera: Aleyrodidae), was collected in Miami-Dade County from gumbo limbo. This was the first report of this insect on the U.S. continent and it is believed to originate from Central America. Since the initial find, there have been numerous other reports, all in Miami-Dade County. It will likely spread to other southern Florida counties.

NOTE: This is not the same whitefly (figus whitefly) that is currently causing defoliation and branch dieback of figus in south Florida.

Host Plants: This whitefly appears to have a very broad host range from palms to woody ornamentals and fruits. Thus far, it has been seen on gumbo limbo (*Bursera simaruba*), *Calophyllum* species, black olive (*Bucida buceras*), copperleaf (*Acalypha wilkesiana*), broadleaf arrowhead (*Sagittaria latifolia*), cocoplum (*Chrysobalanus icaco*), Brazilian pepper (*Schinus terebinthifolius*), wax myrtle (*Myrica cerifera*), live oak (*Quercus virginiana*) and mango (*Mangifera indica*). It has also been reported on several palms which include areca palm (*Dypsis lutescens*), *Veitchia* species, and coconut (*Cocos nucifera*). Additional hosts are likely to be added to the current list.

What are Whiteflies?

They are small, winged insects that belong to the Order Hemiptera which also includes aphids, scales, and mealybugs. These insects typically feed on the underside of leaves with their “needle-like” mouthparts.


Whiteflies can seriously injure host plants by sucking nutrients from the plant causing wilting, yellowing, stunting, leaf drop, or even death. There are more than 75 different whiteflies reported in Florida.

Biology: There is little information on the biology of this new whitefly, however, research is ongoing to describe the life cycle and host range. This whitefly is closely related to giant whitefly, *Aleurodicus*

dugesii, and shares some similarities. Like giant whitefly, the adult is about 3 times larger than other whiteflies that occur in the U.S., and are more docile (slower moving) than other types of whiteflies. The adult whiteflies congregate on the undersides of the leaves to feed and reproduce.



Adult whiteflies



Whitefly eggs

The female whitefly lays her eggs in a spiral pattern on the leaves and also deposits a white, waxy substance on the eggs. The crawler stage hatches from the eggs and crawls around before it starts to feed with its “needle-like” mouthparts. This stage is very small and difficult to see.





Whitefly immature stages

The crawler will molt and go through several immature stages that are oval and initially flat, then more convex. These stages do not resemble a typical insect. Some of these immature stages will secrete long white filaments of wax. It will likely survive year round in south Florida.




Whitefly pupae

Plant Damage: The most noticeable symptoms of an infestation of this whitefly is the abundance of the white, waxy material covering the leaves and also excessive sooty mold. Like other similar insects, these whiteflies will produce “honeydew”, a sugary substance, which causes the growth of sooty mold. The actual effect of an infestation on the health of a plant is unknown; however, whiteflies in general can cause plant decline, defoliation and branch dieback.



Infested palm




Infested gumbo limbo




Sooty mold on black olive

Management in the Landscape: Often, when new pests arrive, they can reach very high populations and can be extremely damaging. It is not uncommon, however, that after several years, the impact is greatly reduced. Although it may be necessary to use insecticides for this pest, it is very important to understand the importance of natural enemies and the need to focus on long-term, biologically based management.

- Monitor plants for early signs of an infestation because it will be easier to manage the pest before it builds to high populations and causes major damage. If you have an infestation on a tree, be sure to search nearby trees as well because this whitefly feeds on many types of trees.
- In the landscape, one parasitoid attacking this whitefly has been identified, *Aleuroctonus vittatus* (Eulophidae), at very high levels. Awareness of natural enemies is very important so they are not also killed while trying to control the whitefly. Protecting natural enemies is a critical component in the long-term control of this pest. Broad spectrum or persistent insecticides often kill a high proportion of predators and parasites, particularly when applied as a foliar spray. Carbamates (i.e. carbaryl), organophosphates (i.e. malathion) and pyrethroids (i.e. bifenthrin, cyfluthrin) are especially toxic to natural enemies.



Parasitized whitefly pupae



Parasitoid

- You may achieve excellent control with one of the methods listed below, but remember the white, waxy material and the sooty mold on the plant will take time to wear off unless physically washed off.
- Washing plants off with water can be an effective tool to help manage whiteflies for small infestations or small plants. But, for it to be effective, you must remove the immature stages and eggs from the leaves with the wash.
- Using a **horticultural oil** or **insecticidal soap** can also help control this pest. These types of products are strictly contact so thorough coverage of the infested leaves is required. Typically, several applications are required 7-10 days apart. Be careful about using these types of products under high temperatures because they can cause damage to plants.
- If the infestation is large, an insecticide may be needed to control the whitefly population. It is extremely important to use the appropriate insecticides, methods, and timing in order to get the best control with the least amount of detriment to the natural enemies or the environment. There are several insecticide options for both professional use (Table 1) and homeowner use (Table 2). Many of the insecticides for professional use are available in more than one formulation (i.e. wettable powder, liquid, soluble granules, granules, pellets) so you can choose the best fit for your situation.
- Contact insecticides are typically sprayed on the foliage or other infested parts of the plant or in the soil for soil-dwelling insects. Depending on the insecticide, either the insect must come into contact with the insecticide or must feed on the plant with the insecticide. Spray coverage must be thorough to get the best results, particularly in cases like this when the insect is primarily on the underside of the leaves. In general, foliar sprays are active for a few weeks and usually require more than one application. However, some of these products can be very useful for quick knockdown which can be very important with bad infestations.
- A systemic insecticide can be applied directly to the infested plant or to the soil. Soil applications include drenching the soil, spreading a granular formulation, or burying a pellet. Some products can also be applied as a basal trunk spray or injection into the trunk. Systemic insecticides can also be sprayed on the foliage, but often provide longer control when applied to the soil or trunk. However, it is not recommended to use the same insecticide (active ingredient) on the leaves that you use in the soil or on the trunk.
- Misuse or overuse of any insecticide can cause problems such as insecticide resistance, secondary pest problems, environmental contamination, and detrimental effects on non-target organisms. The **site** and **method of application** must be on the insecticide label. Always follow the label directions – **“The label is the law”**. If it is necessary for you to apply several applications of insecticides, it is recommended to rotate among different chemical classes .

Table 1. Insecticides labeled for professional use against whiteflies in Florida

Active Ingredient	Chemical Class	Trade Name(s)	Contact or Systemic
Abamectin	Avermectins	Avid	Contact
Acephate	Organophosphate	Orthene	Systemic
Acetamiprid	Neonicotinoid	TriStar	Systemic
Azadirachtin	Botanical	Azatin; Azatrol	Contact
<i>Beauveria bassiana</i>	Microbial	Botanigard	Contact
Bifenthrin	Pyrethroid	Bifenthrin Pro; Onyx; Talstar	Contact
Buprofezin	IGR	Talus*	Contact
Carbaryl	Carbamate	Sevin	Contact
Clothianidin	Neonicotinoid	Arena; Aloft**	Systemic
Cyfluthrin	Pyrethroid	Tempo; Decathlon*	Contact
Cypermethrin	Pyrethroid	Demand	Contact
Deltamethrin	Pyrethroid	DeltaGard	Contact
Diflubenzuron	IGR	Adept*; Dimilin*	Contact
Dinotefuran	Neonicotinoid	Safari	Systemic
Fenoxycarb	IGR	Preclude	Contact
Fenpropathrin	Pyrethroid	Tame	Contact
Fonicamid	Antifeedant	Aria*	Contact
Fluvalinate	Pyrethroid	Mavrik; Aquaflow	Contact
Imidacloprid	Neonicotinoid	Marathon*; Merit; CoreTect; Discus* **; Allectus**	Systemic
Lambda-cyhalothrin	Pyrethroid	Scimitar	Contact
Malathion	OP	Malathion	Contact
Novaluron	IGR	Pedestal*	Contact
Paraffinic Oil	Oil	Horticultural Oil	Contact
Soap	Soap	Insecticidal Soap	Contact
Pymetrozine	Antifeedant	Endeavor	Contact
Pyridaben	Acaricide	Sanmite	Contact
Pyriproxyfen	IGR	Distance	Contact
S-Kinoprene	IGR	Enstar II*	Contact
Spinosad	Microbial	Conserve	Contact
Spiromesifen	IGR	Forbid 4F; Judo*	Contact
Thiamethoxam	Neonicotinoid	Flagship; Meridian	Systemic

* For production nursery, greenhouse and/or interiorscape use only.

**Product also contains a pyrethroid

Table 2. Insecticides labeled for homeowner use against whiteflies in Florida

Active Ingredient	Chemical Class	Trade Name(s)
Bifenthrin	Pyrethroid	Ortho Bug-B-Gon Max Lawn & Garden Insect Killer
Carbaryl	Carbamate	Sevin
Cyfluthrin	Pyrethroid	Bayer Advanced Rose & Flower Insect Killer; Schultz Lawn & Garden Insect Killer
Dinotefuran	Neonicotinoid	Green Light Tree & Shrub Insect Control with Safari
Imidacloprid	Neonicotinoid	Bayer Advanced Lawn Complete Insect Killer; Bayer Advanced Tree & Shrub Insect Control ; Ortho Max
Lambda-cyhalothrin	Pyrethroid	Spectracide Triazicide Once & Done Insect Killer
Malathion	Organophosphate	Green Light Malathion; Ortho Malathion Plus Insect Spray
Neem oil	Botanical	Bonide Safer BioNeem; Green Light Neem; Green Light Rose Defense; Southern Ag Triple Action Neem Oil
Paraffinic oil	Biorational	Sun Spray Horticultural Oil
Permethrin	Pyrethroid	Hi-Yield Indoor/Outdoor Broad Use Insecticide
Potassium salts	Biorational	Safer's Insecticidal Soap
Pyrethrins	Botanical	Bonide Yard & Garden Insect Killer; Spectracide Rose & Flower Insect Spray

For more information, contact your local Extension agent for additional information or <http://trec.ifas.ufl.edu/mannion/>

Catharine Mannion, Ph.D., Associate Professor and Extension Specialist
University of Florida, IFAS, Tropical Research and Education Center, Homestead, FL

Photo Credits: H. Glenn, UF/IFAS, Tropical Research and Education Center, Homestead, FL

The use of trade names in this publication is solely for the purpose of providing specific information. UF/IFAS does not guarantee or warranty the products named, and references to them in this publication does not signify our approval to the exclusion of other products of suitable composition. All chemicals should be used in accordance with directions on the manufacturer's label. Check the label carefully as this information does change and our publications may not have the latest information. Use pesticides safely. Read and follow directions on the manufacturer's label.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other extension publications, contact your county Cooperative Extension service.

U.S. Department of Agriculture, Cooperative Extension Service, University of Florida, IFAS, Florida A. & M. University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Larry Arrington, Dean.